

Review Article

Chronic pain, Sodium channel and Gene mutationDr Santanu Mallik DA^{*1}, Dr V. G Sawant²¹Department of Anatomy Dr D Y Patil Medical College, Navi Mumbai, India²Head & Professor, Department of Anatomy, Dr D Y Patil Medical College, Navi Mumbai, India***Corresponding author**

Dr Santanu Mallik DA

Email: drmallik@rediffmail.com

Abstract: Anatomy of human pain pathway is complex. It starts with the primary sensory neurons in the periphery, conduction to the spinal cord through afferent neurons, and processing at multiple higher levels that include the dorsal horn, spinal projections, thalamus, and cortex. Transmission of painful stimulus and its processing involves different ion channels, receptors, and various chemical mediators. Different sodium channels, especially Nav1.7 are expressed preferentially in most slowly conducting nociceptive neurons and in sympathetic neurons. Dysregulated expression of voltage-gated sodium channels can produce neuronal hyperexcitability associated with severe pain, whereas loss of the Nav1.7 channel in patients leads to indifference to pain. Interestingly, certain factors of human pain perception can be inherited like acute pain thresholds, efficacy of analgesics, inclination of developing chronic pain, etc. Mutation of the genes encoding for ion channels can cause heritable pain disorders by abnormal channel function or expression. Discovery of genetic mutations and of additional variants that likely to be identified in the future will open up the possibility of understanding chronic pain accurately, at the molecular level. The understanding of the pain pathway at the molecular level may lead to new therapeutic strategies that will bring us closer to more effective treatments for pain.

Keywords: Chronic pain, Sodium channel (Nav), Dorsal root ganglion, inherited pain disorders, inherited erythromelalgia

INTRODUCTION

Sodium channels are integral membrane proteins that form ion channels, conducting sodium ions (Na^+) through a cell's plasma membrane present in excitable cells such as neurons, myocytes, responsible for the rising phase of action potentials[1,2]. Pain pathways start with primary sensory neurons of dorsal root ganglion and trigeminal neurons. The peripheral stimulus that develops pain perception depends on the presence of functional voltage-gated sodium channels. Several studies shown that, after injury to their axons, neurons can display changes in excitability due to increased sodium channel expression. Though studies showed that, in some pain syndromes, hyperexcitability or increased baseline sensitivity of these cells leads to abnormal bursting that can produce chronic pain but the underlying molecular mechanisms are not fully understood [3-5].

Changes in Dorsal root ganglion after injury

Early studies [6,7] described that, after injury to their axons, motor neurons show changes in excitability, suggesting increased sodium channel expression over the cell body and the dendrites, and subsequently similar changes were observed in sensory neurons[8,9]. Abnormal sodium channel aggregation at the tips of injured axons also has been observed[10-12]

that abnormally increases in sodium conductance which can lead to inappropriate, repetitive firing[13-15] causing chronic pain. These observations, together with experimental and clinical observations on partial efficacy of sodium channel blocking agents in neuropathic pain, established a link between sodium channel activity and sensory neuron hyperexcitability producing pain [16-19].

Sodium Channels in Sensory Neurons

Over the past decade, a family of nine structurally related Voltage-gated sodium channels is encoded within mammals by different genes. Dorsal root Ganglion (DRG) neurons, which had been known to display multiple, distinct sodium currents [20-23] express at least six sodium channel transcripts [24]. Out of nine pore-forming voltage-gated sodium channel (Nav 1.1 to Nav 1.9), Sodium Channel Nav 1.7 and Nav 1.8 are important in nociceptive pathway. Nav 1.7 has been detected in DRG (A-Beta and C fibers), trigeminal ganglion and sympathetic ganglion neuron (superior cervical ganglion). Its presence is less in CNS, smooth myocytes [25].

Sodium Channel Gene Expression Is Altered After Injury to DRG Neurons

Waxman *et al* observed that in addition to production of excess channels, there is a switch in the type of channels produced cause a significant up-regulation of expression of the previously silent sodium channel gene ((SNSyPN3, NaN, PN1, and NaG) in DRG neurons after axotomy. This finding was followed by demonstration of down-regulation of the gene expression, which can persist as long as 210 days [26], which is consistent with the long-lasting changes in gene expression that have been described in these cells [27]. These changes may poise DRG neurons to fire spontaneously, or at inappropriately high frequencies, after injury. Increased sodium channel densities, in themselves, will tend to lower threshold [14].

Sodium Channel Expression in Inflammatory Pain Models

Several studies have demonstrated that inflammatory molecules such as prostaglandins and serotonin can modulate sodium currents in DRG neurons [29], possibly acting through a cyclic AMP-protein cascade [30]. However, whether sodium channel gene expression is affected in inflammatory models of pain had not been addressed. But some studies have shown a persistent increase in sodium channel in DRG neurons within 24 hrs of injection of carageenan[31] in rat . The mechanism responsible for this inflammation-associated change in sodium channel expression is not known. Interestingly, Nerve Growth Factor (NGF) normally is produced in peripheral target tissues by supporting cells that include fibroblasts, Schwann cells, and NGF production is stimulated in immune cells, and increased NGF levels have been observed in the local area, raising the possibility that inflammation may indirectly trigger changes in sodium channel gene expression via changes in neurotrophin levels [32,33].

Now it is clear to us that, dysregulated expression of several sodium channels has been associated with chronic pain induced by trauma, inflammation or metabolic disorders (e.g. diabetes) and blocking of sodium channels can ameliorate pain symptoms [34-37]. However, one particular sodium channel isoform Nav1.7, has emerged as a major focus in studies on pain, in large part because recent studies have identified Nav1.7 as a key contributor to nociceptive neuronal excitability, which is linked to different human pain disorders [38-40].

Certain factors of human pain experience are found to be inherited like acute pain thresholds, efficacy of analgesics. Mutation of the genes encoding for ion channels can cause heritable pain disorders by abnormal channel function or expression. These diseases caused by these gene mutations are rare but refractory to conventional treatment.

Contribution of Nav1.7 to pain: human studies

Inherited sodium channelopathies underlie several neurological and muscular diseases [41-44]. In 2004 a painful syndrome erythromelalgia (also called erythromalgia) was found in two Chinese families with two independent mutations in SCN9A, the gene that encodes Nav1.7 [45]. Since 2004, several more mutations in SCN9A have been linked to inherited erythromelalgia [46]. Additionally, other mutations in SCN9A have been linked to inherited paroxysmal extreme pain disorder [47] and, more recently, individuals with complete loss of functional Nav1.7 have been reported to be 'indifferent' to pain [48,49].

Inherited erythromelalgia

Life-long symptoms of early-onset (as early as 1 year-old) are characterized by episodes of burning pain triggered by mild warmth or exercise, together with erythema and mild swelling in the hands and feet and sometimes in the ears or face [50-53]. The frequency and severity of pain episodes increase with age, with each episode lasting minutes to hours. Typically, patients with early-onset inherited erythromelalgia do not report autonomic abnormalities, such as orthostatic hypotension or gastrointestinal symptoms. Partial relief of symptoms comes from cooling the affected extremities. Linkage analysis identified the disease locus on chromosome 2 (2q31–32) [54] where a cluster of voltage-gated sodium-channel genes, including SCN9A, is known to exist and, subsequently, the molecular target was identified as Nav1.7[55]. Indeed, two independent mutations (I848T and L858F) in two sporadic early-onset erythromelalgia cases from China [55,56], which have also been reported from multigeneration French (I848T) and Canadian (L858F) families[54]. Familial adult-onset erythromelalgia has been described recently in a family from the USA in which coding mutations of Nav1.7 were excluded as the causative factor [57]. Thus, it is possible that mutations in noncoding regions of the SCN9A gene that cause increased expression of the channel or indeed other target genes might cause inherited erythromelalgia.

Paroxysmal extreme pain disorder (PEPD)

A different set of mutations in Nav1.7 has been associated with another autosomal dominant painful disorder, PEPD[40], which was known as familial rectal pain previously [58,59]. PEPD is characterized by life-long pain episodes, accompanied by tonic posturing and immediately followed by flushing of the lower limbs, in a uni- or bi-lateral fashion, which start within the first days after birth and are triggered by defecation or probing of the anal or genital areas. The frequency of the rectal-pain episodes decreases with age; ocular and mandibular pain are sometimes triggered by cold or irritants, becoming more prominent complaints. In contrast to inherited erythromelalgia, which is generally refractory to pharmacological treatment, PEPD patients respond to

treatment with carbamazepine with almost complete cessation of spontaneous or induced attacks [40,58]. This patient has more severe symptoms compared with his father and with patients in another family carrying the R996C mutation alone [60].

Congenital indifference to pain (CIP)

Patients with congenital insensitivity (or indifference) to pain (CIP) show varying degrees of deficits in sensing, perceiving and reacting to painful stimuli but their other sensory modalities are intact [61,62]. The designation of patients as 'insensitive' or 'indifferent' to pain is a subject of controversy that has led to the use of the term channelopathy-associated CIP for a subset of these patients with null mutations in Nav1.7. Typically, these patients present with a history of not experiencing any form of pain anywhere on their bodies, even after burns, bone fractures or severe injuries to their lips and tongues, and they do not experience visceral pain [63-65]. Despite the fact that Nav1.7 is produced within sympathetic ganglion neurons [66-70], patients with channelopathy-associated CIP do not show apparent deficits in autonomic function, such as whole-body thermoregulation or cardiac or respiratory rhythm, and have a normal axon-reflex response to histamine [48,63]. Interestingly, channelopathy-associated CIP patients have deficits in the sense of smell [48]. Three loss-of-function mutations in SCN9A were identified initially in unrelated consanguineous Pakistani families [63]. Subsequently several other mutations, including deletions in coding sequence that truncate the protein and deletion of intronic sequences that suggest splicing defects, were identified in ethnically diverse patients [64]. Surprisingly, these cases included patients who were born into families that are apparently non-consanguineous, which suggests a de novo mutation of one of the alleles and the inheritance of the second mutant allele from one of the parents, who would be an asymptomatic carrier. This observation suggests that SCN9A-related CIP might underlie some cases of idiopathic CIP in non-consanguineous families [62].

Familial hemiplegic migraine (FHM)

It is a form of migraine headache that runs in families. It is inherited in autosomal dominant fashion, and aura symptoms may be associated with moderate to severe motor weakness, ataxia, and seizures. Mutations in the CACNA1A (on chromosome 19p3 coding for calcium channel Cav2.1) [71,72], ATP1A2 (encoding the Na⁺/K⁺ ATPase α 2 subunit leads to impaired pump function there by leading to loss of ion gradient at channels), SCN1A gene mutation is in the highly conserved region of the sodium channel responsible for fast inactivation [73], and PRRT2 genes are found to cause familial hemiplegic migraine. The function of the protein produced from the PRRT2 gene is unknown although it interacts with a protein that helps control interneuronal signaling.

Familial episodic pain syndrome (FEPS)

It is associated with episodes of upper body pain of severe intensity. The pain is triggered by fasting, physical stress, or fatigue. It was first found in a Colombian family and first disease linked with transient receptor potential channels (TRP channels). These are a group of ion channels located mostly on the plasma membrane of numerous human cell types and mediate a variety of sensations like pain, hotness, warmth, coldness, taste, pressure, and vision. A nonsense mutation in TRPA1 gene is found to be linked to the syndrome [74].

Irritable Bowel Syndrome (IBS)

Several studies suggest a strong genetic component causing familial IBS but association of the candidate genes are yet to be found [75].

CONCLUSION

Sodium channels are significant participants in electro-genesis within sensory neurons, including DRG and trigeminal nerves and they probably subserve multiple functions (transduction, signal amplification, action potential electrogenesis, etc.) and interact in a complex manner in trauma, inflammation or metabolic disorders (e.g. diabetes). Mutation of the genes encoding for ion channels can cause heritable pain disorders by abnormal channel function or expression. It is quite likely, in our opinion, that sodium channel blockade will emerge as a viable strategy for pharmacologic treatment of pain but require further careful studies. Discovery of genetic mutations and of additional variants that will likely be identified in the future open up the possibility of understanding chronic pain more fully, at the molecular level.

REFERENCES

1. Jessell TM, Kandel ER, Schwartz JH; Principles of Neural Science (4th ed.). New York: McGraw-Hill, 2000; 154-169.
2. Bertil Hille; Ion Channels of Excitable Membranes (3rd ed.). Sunderland, Mass: Sinauer, 2001; 73-77.
3. Ochoa J, Torebjork HE; Paraesthesiae from ectopic impulse generation in human sensory nerves. *Brain*, 1980; 103: 835-854.
4. Nordin M, Nystrom B, Wallin U, Hagbarth KE; Ectopic sensory discharges and paresthesiae in patients with disorders of peripheral nerves, dorsal roots and dorsal columns. *Pain*, 1984; 20(3), 231-245.
5. Devor M; in Textbook of Pain, eds Wall P D, Melzack R Churchill Livingstone, Edinburgh, 1994; 79-101.
6. Eccles JC, Libet B, Young RR; The behaviour of chromatolysed motoneurons studied by intracellular recording. *The Journal of physiology*, 1958; 143(1):11-40.
7. Kuno M, Llinas R; Enhancement of synaptic transmission by dendritic potentials in

- chromatolysed motoneurons of the cat. The Journal of physiology, 1970; 210(4):807-821.
8. Gallego R, Ivorra I, Morales A; Effects of central or peripheral axotomy on membrane properties of sensory neurons in the petrosal ganglion of the cat. The Journal of Physiology, 1987; 391(1):39-56.
9. Gurtu S, Smith PA; Electrophysiological characteristics of hamster dorsal root ganglion cells and their response to axotomy. Journal of neurophysiology, 1988;59(2):408-423.
10. Devor M, Keller CH, Deerinck TJ, Levinson SR, Ellisman MH; Na⁺ channel accumulation on axolemma of afferent endings in nerve end neuromas in Apterionotus. Neuroscience letters, 1989;102(2):149-154.
11. England JD, Gamboni F, Ferguson MA, Levinson SR; Sodium channels accumulate at the tips of injured axons. Muscle & nerve, 1994; 17(6):593-598.
12. England JD, Happel LT, Kline DG, Gamboni F, Thouron CL, Liu ZP, Levinson SR; Sodium channel accumulation in humans with painful neuromas. Neurology, 1996; 47(1):272-276.
13. Waxman SG, Ritchie JM; Organization of ion channels in the myelinated nerve fiber. Science, 1985; 228(4707):1502-1507.
14. Matzner O, Devor M; Na⁺ conductance and the threshold for repetitive neuronal firing. Brain research, 1992; 597(1):92-98.
15. Matzner O, Devor M; Hyperexcitability at sites of nerve injury depends on voltage-sensitive Na⁺ channels. Journal of Neurophysiology, 1994;72(1):349-359.
16. Chabal C, Russell LC, Burchiel KJ; The effect of intravenous lidocaine, tocainide, and mexiletine on spontaneously active fibers originating in rat sciatic neuromas. Pain, 1989; 38(3):333-338.
17. Devor M, Wall PD, Catalan N; Systemic lidocaine silences ectopic neuroma and DRG discharge without blocking nerve conduction. Pain, 1992;48(2):261-268.
18. Omana-Zapata I, Khabbaz MA, Hunter JC, Bley KR; QX-314 inhibits ectopic nerve activity associated with neuropathic pain. Brain research, 1997; 771(2):228-237.
19. Rizzo MA; Successful treatment of painful traumatic mononeuropathy with carbamazepine: insights into a possible molecular pain mechanism. Journal of the neurological sciences, 1997;152(1):103-106.
20. Kostyuk PG, Veselovsky NS, Tsyandryenko AY; Ionic currents in the somatic membrane of rat dorsal root ganglion neurons I: Sodium currents Neuroscience, 1981; 6: 2423-2430.
21. Roy ML, Narahashi T; Differential properties of tetrodotoxin-sensitive and tetrodotoxin-resistant sodium channels in rat dorsal root ganglion neurons. The Journal of neuroscience, 1992;12(6):2104-2111.
22. Caffrey JM, Eng DL, Black JA, Waxman SG, Kocsis JD; Three types of sodium channels in adult rat dorsal root ganglion neurons. Brain research, 1992;592(1):283-297.
23. Elliott AA, Elliott JR; Characterization of TTX-sensitive and TTX-resistant sodium currents in small cells from adult rat dorsal root ganglia. The Journal of Physiology, 1993; 463:39.
24. Black JA, Dib-Hajj S, Baker D, Newcombe J, Cuzner ML, Waxman SG; Sensory neuron-specific sodium channel SNS is abnormally expressed in the brains of mice with experimental allergic encephalomyelitis and humans with multiple sclerosis. Proceedings of the National Academy of Sciences, 2000; 97(21):11598-11602.
25. Cummins TR, Sheets PL, Waxman SG; The roles of sodium channels in nociception: Implications for mechanisms of pain. Pain, 2007; 131: 243-257.
26. Waxman SG, Dib-Hajj S, Cummins TR, Black JA; Sodium channels and pain. Proceedings of the National Academy of Sciences, 1999; 96(14):7635-7639.
27. Dib-Hajj S, Black JA, Felts P, Waxman SG; Down-regulation of transcripts for Na channel α -SNS in spinal sensory neurons following axotomy. Proceedings of the National Academy of Sciences, 1996; 93(25):14950-14954.
28. Matzner O, Devor M; Na⁺ conductance and the threshold for repetitive neuronal firing. Brain research, 1992;597(1):92-98.
29. Gold MS, Reichling DB, Shuster MJ, Levine JD; Hyperalgesic agents increase a tetrodotoxin-resistant Na⁺ current in nociceptors. Proceedings of the National Academy of Sciences, 1996; 93(3):1108-1112.
30. England S, Bevan S, Docherty RJ; PGE2 modulates the tetrodotoxin-resistant sodium current in neonatal rat dorsal root ganglion neurons via the cyclic AMP-protein kinase A cascade. The Journal of Physiology, 1996;495(Pt 2):429.
31. Gould HJ, England JD, Liu ZP, Levinson SR; Rapid sodium channel augmentation in response to inflammation induced by complete Freund's adjuvant. Brain research, 1998; 802(1): 69-74.
32. Woolf CJ, Safieh-Garabedian B, Ma QP, Crilly P, Winter J; Nerve growth factor contributes to the generation of inflammatory sensory hypersensitivity. Neuroscience, 1994; 62(2):327-331.
33. Weskamp G, Otten U; An enzyme-linked immunoassay for nerve growth factor (NGF): a tool for studying regulatory mechanisms involved in NGF production in brain and in peripheral tissues. Journal of neurochemistry, 1987;48(6):1779-1786.
34. Waxman SG; Transcriptional channelopathies: an emerging class of disorders. Nat. Rev. Neurosci., 2001; 2: 652-659.
35. Waxman SG, Hains BC; Fire and phantoms after spinal cord injury:Na⁺ channels and central pain. Trends Neurosci., 2006; 29: 207-215

36. Wood JN, Boorman JP, Okuse K, Baker MD; Voltage-gated sodium channels and pain pathways. *J. Neurobiol*, 2004; 61: 55–71.
37. Lai J, Porreca F, Hunter JC, Gold MS; Voltage-gated sodium channels and hyperalgesia. *Annu. Rev. Pharmacol. Toxicol*, 2004; 44: 371–397.
38. Waxman SG, Dib-Hajj S; Erythralgia: molecular basis for an inherited pain syndrome. *Trends Mol. Med.*, 2005; 11: 555–562.
39. Cox JJ, Reimann F, Nicholas AK, Thornton G, Roberts E, Springell K, Woods CG, et al.; An SCN9A channelopathy causes congenital inability to experience pain. *Nature*, 2006; 444: 894–898.
40. Fertleman CR, Baker MD, Parker KA, Moffatt S, Elmslie FV, Abrahamsen B, Rees M, et al.; SCN9A mutations in paroxysmal extreme pain disorder: allelic variants underlie distinct channel defects and phenotypes. *Neuron*, 2006; 52: 767–774.
41. Meisler MH, Kearney JA; Sodium channel mutations in epilepsy and other neurological disorders. *J. Clin. Invest*, 2005; 115: 2010–2017.
42. George AL; Inherited disorders of voltage-gated sodium channels. *J. Clin. Invest*, 2005; 115, 1990–1999.
43. Moss AJ, Kass RS; Long QT syndrome: from channels to cardiac arrhythmias. *J. Clin. Invest*, 2005; 115: 2018–2024.
44. Cannon SC; Pathomechanisms in channelopathies of skeletal muscle and brain. *Annu. Rev. Neurosci.*, 2006; 29: 387–415.
45. Yang Y, Wang Y, Li S, Xu Z, Li H, Ma L, Shen Y, et al.; Mutations in SCN9A, encoding a sodium channel alpha subunit, in patients with primary erythralgia. *J. Med. Genet*, 2004; 41: 171–174.
46. Dib-Hajj SD, Rush AM, Cummins TR, Waxman SG; Mutations in the sodium channel Nav1.7 underlie inherited erythralgia, 2006; 3: 343–350.
47. Fertleman CR, Baker MD, Parker KA, Moffatt S, Elmslie FV, Abrahamsen B, Rees M, et al.; SCN9A mutations in paroxysmal extreme pain disorder: allelic variants underlie distinct channel defects and phenotypes. *Neuron*, 2006; 52: 767–774.
48. Goldberg YP, MacFarlane J, MacDonald ML, Thompson J, Dube MP, Mattice M, Hayden MR; Loss-of-function mutations in the Na(v)1.7 gene underlie congenital indifference to pain in multiple human populations, 2007; 71: 311–319.
49. Ahmad S, Dahllund L, Eriksson AB, Hellgren D, Karlsson U, Lund PE, Krupp JJ, et al.; A stop codon mutation in SCN9A causes lack of pain sensation. *Hum. Mol. Genet.*, 2007; 16: 2114–2121.
50. Novella SP, Hisama FM, Dib-Hajj SD, Waxman SG; A case of inherited erythralgia. *Nat Clin Pract Neurol*, 2007; 3: 229–234.
51. Hisama FM, Dib-Hajj SD, Waxman SG; SCN9A-related inherited erythralgia. In *GeneReviews* at GeneTests: Medical Genetics Information Resource 2013.
52. Drenth JP, Michiels JJ; Clinical characteristics and pathophysiology of erythralgia and erythralgia. *Am. J. Med*, 1992; 111–114.
53. van Genderen PJ, Michiels JJ, Drenth JP; Hereditary erythralgia and acquired erythralgia. *Am. J. Med. Genet*, 1993; 45: 530–532.
54. Drenth JP, Finley WH, Breedveld GJ, Testers L, Michiels JJ, Guillet G, Heutink P; The primary erythralgia-susceptibility gene is located on chromosome 2q31–32. *Am. J. Hum. Genet*, 2001; 68: 1277–1282.
55. Yang Y, Wang Y, Li S, Xu Z, Li H, Ma L, Shen Y; Mutations in SCN9A, encoding a sodium channel alpha subunit, in patients with primary erythralgia. *J. Med. Genet*, 2004; 41: 171–174.
56. Han C, Rush AM, Dib-Hajj SD, Li S, Xu Z, Wang Y, Waxman SG; Sporadic onset of erythralgia: a gain-of-function mutation in Nav1.7. *Ann. Neurol*, 2006; 59: 553–558.
57. Burns TM, Te Morsche RHM, Jansen JBMJ, H Drenth JP; Genetic heterogeneity and exclusion of a modifying locus at 2q in a family with autosomal dominant primary erythralgia. *Br. J. Dermatol*, 2005; 153: 174–177.
58. Bednarek N, Arbuès AS, Motte J, Sabouraud P, Plouin P, Morville P; Familial rectal pain: a familial autonomic disorder as a cause of paroxysmal attacks in the newborn baby. *Epileptic Disord*, 2005; 7: 360–362.
59. Dugan RE; Familial rectal pain. *Lancet* 1, 854. Schubert R. and Cracco, JB; Familial rectal pain: a type of reflex epilepsy? *Ann. Neurol*, 1972; 32: 824–826.
60. Schubert R, Cracco JB; Familial rectal pain: a type of reflex epilepsy? *Ann. Neurol*, 1992; 32: 824–826.
61. Verhoeven K, Timmerman V, Mauko B, Pieber TR, De Jonghe P, Auer-Grumbach M; Recent advances in hereditary sensory and autonomic neuropathies. *Curr. Opin. Neurol*, 2006; 19: 474–480.
62. Nagasako EM, Oaklander AL, Dworkin RH; Congenital insensitivity to pain: an update, 2003; 101: 213–219.
63. Cox JJ, Reimann F, Nicholas AK, Thornton G, Roberts E, Springell K, Woods CG; An SCN9A channelopathy causes congenital inability to experience pain. *Nature* 444, 894–898 inability to experience pain, 2006; 444: 894–898.
64. Goldberg YP, MacFarlane J, MacDonald ML, Thompson J, Dube MP, Mattice M, Hayden MR; Loss-of-function mutations in the Na(v)1.7 gene underlie congenital indifference to pain in multiple human populations, 2007; 71: 311–319.
65. Ahmad S, Dahllund L, Eriksson AB, Hellgren D, Karlsson U, Lund PE, Krupp JJ; A stop codon

- mutation in SCN9A causes lack of pain sensation. Hum. Mol. Genet, 2007; 16: 2114–2121.
66. Black JA, Dib-Hajj S, McNabola K, Jeste S, Rizzo MA, Kocsis JD, Waxman SG; Spinal sensory neurons express multiple sodium channel alpha-subunit mRNAs. Brain Res, 1996; 43: 117–131.
67. Toledo-Aral JJ, Moss BL, He ZJ, Koszowski AG, Whisenand T, Levinson SR, Mandel G; Identification of PN1, a predominant voltage-dependent sodium channel expressed principally in peripheral neurons. Proc. Natl. Acad. Sci. U. S. A., 1997; 94: 1527–1532.
68. Sangameswaran L, Fish LM, Koch BD, Rabert DK, Delgado SG, Ilnicka M, Hunter JC; A novel tetrodotoxin-sensitive, voltage-gated sodium channel expressed in rat and human dorsal root ganglia. J. Biol. Chem, 1997; 272: 14805–14809.
69. Felts PA, Yokoyama S, Dib-Hajj S, Black JA, Waxman SG; Sodium channel alpha-subunit mRNAs I, II, III, NaG, Na6 and HNE (PN1) – different expression patterns in developing rat nervous system. Brain Res., 1997; 45: 71–82.
70. Rush AM, Dib-Hajj SD, Liu S, Cummins TR, Black JA, Waxman SG; A single sodium channel mutation produces hyper- or hypoexcitability in different types of neurons. Proc. Natl. Acad. Sci. U. S. A., 2006; 103: 8245–8250.
71. Joutel A, Bousser MG, Biousse V, Labauge P, Chabriat H, Nibbio A, et al.; A gene for familial hemiplegic migraine maps to chromosome 19. Nat Genet, 1993; 5: 40-45.
72. Dichgans M, Freilinger T, Eckstein G, Babini E, Lorenz-Depiereux B, Biskup S, et al.; Mutation in the neuronal voltage-gated sodium channel SCN1A in familial hemiplegic migraine. Lancet, 2005; 366: 371-377.
73. Heron SE, Dibbens LM; Role of PRRT2 in common paroxysmal neurological disorders a gene with remarkable pleiotropy. J Med Genet, 2013; 50: 133-139.
74. Kremeyer B, Lopera F, Cox JJ, Momin A, Rugiero F, Marsh S, et al.; A gain-of-function mutation in TRPA1 causes familial episodic pain syndrome. Neuron, 2010; 66: 671-680.
75. Saito YA, Talley NJ; Genetics of irritable bowel syndrome. Am J Gastroenterol, 2008; 103: 2100-2104.